



PRODUCTION OF TRUE-TO-TYPE GUAVA PLANTS THROUGH CLONAL PROPAGATION

Nitin Soni¹, S.K. Pandey², S. S. Singh³, S.R.K. Singh⁴, A. Mishra⁵, S.S. Baghel⁶ and Pavan Kumar Kaurav⁷

¹RA-NICRA, ICAR- ATARI, Zone VII, JNKVV campus, Jabalpur.(MP) Pin 482004

²Professor, Department of Horticulture College of Agriculture, JNKVV, Jabalpur (MP) Pin 482004

³Assistant Professor, MGCGVV, Chitrakoot, Dist Satna (MP)

⁴Senior Scientist, ICAR- ATARI, Zone VII, JNKVV campus, Jabalpur (MP) Pin 482004

⁵Director, ICAR- ATARI, Zone VII, JNKVV campus, Jabalpur (MP) Pin 482004

⁶College of Agriculture, Rewa (MP)

⁷Team Leader, MPWSRP, Keolari, Seoni (MP)

ABSTRACT

Different concentrations of IBA (0, 1000, 2000 and 3000 ppm) on guava semi hard wood and hardwood cuttings planted in different rooting media (Soil, Silt, Silt +Soil, Soil + Silt+ FYM, Soil+ Silt +Vermi compost) under net house condition poly bags. Significantly higher success percentage (33.09, 36.63%) was noted in both of cuttings treated with 3000 ppm IBA as compared to control (21.68 and 23.9 %) in semihard wood and hard wood respectively. The plants produced 5.44 and 5.33 numbers of leaves, 14.37 and 10.91 cm shoot length and 26.58 and 24.10 number of roots as compared to control . The study provided useful information on guava clonal multiplication through cutting and IBA impact on samihard wood and hard wood cuttings.

Keywords: *Psidium guajava*; IBA; protected environment; asexual; propagation technique; cuttings

I. INTRODUCTION

Guava (*Psidium guajava* L.) the poor man's fruit or apple of the tropics is popular in tropical and subtropical climates. It is native to tropical America stretching from Mexico to Peru (Jaiswal *et.al.*(1987) and Martin *et. al.* (1987)). Guava is cultivated in every tropical and subtropical country around the world (Smart, D. R. *et.al.* , 2001). This is a delicious fruit and is very nutritious and exceptionally rich in ascorbic acid and several minerals useful for human health (Wilson *et.al.* , 1980). Besides its high nutritional value, it bears heavy crop every year and gives good economic returns (Singh *et. al.*, 2000).

Guava propagation through seed does not produce true-to-type plants while clonal propagation has assured true-to-type plants. Maqbool and Khan *et. al.* (1973) reported that guava is commercially propagated from seeds in Pakistan. Guava, if propagated through seed, exhibits a great variation due to inevitable heterozygosity. Moreover, seed propagated plants come into bearing much later than vegetatively propagated plants. Through seed propagation unique characters of a certain variety cannot be preserved or multiplied. Seed propagation does not permit the utilization of superior important characters of a certain rootstock such as disease tolerance (viral fungal or bacterial), adaptability to varying agro-ecological conditions, manipulation of tree growth (dwarfness) and better influence of certain rootstock. Vegetative propagation is, therefore, inevitable in guava Urooba, P. *et.al.* (2008) . In fruit trees, several

vegetative propagation techniques as air layering, root cuttings and stooling, have been tried with varying success rate to increase productivity and gains by clonal propagation and selection (Morton, J. (1987) , Pathak, R.K. and R. Dwivedi. 1996)). However, these techniques are still not commercially viable due to varying rate of success, absence of tap root system and cumbersome process. Healthy planting material is essential to achieve good yield and quality produce. A well-established commercial nursery must improve the way of producing planting material using modern technology as there is the potential to produce true-to-type guava nursery plants with soft wood cuttings Abdullah *et.al.*(2006) In Punjab, guava is generally propagated from seeds and the seedlings are variable in both plant and fruit characteristics. Establishment of orchard through seedlings is not recommended at present time; as most of these seedlings will not be like the parental type in yield, taste and fruit flesh color.

The major issue in guava plantation is discriminate multiplication of plants from unreliable sources by nurserymen (Singh *et.al.*, 2005). Non-availability of quality planting material and consequent substitution of poor quality seedlings have adversely affected the guava production. True-to-type initial planting material is basic need to ensure both quality and quantity in guava (Singh *et.al.*, 2005). Breeding programmes for perennial plants like fruit trees are time consuming because of their slow growth rate and long generation time. In present context, rapid methods of propagation become very important when planting material is limited due to scarcity of a clone or varieties or due to sudden expansion in acreage. Adventitious root formation is a key component of clonal propagation of selected woody plants Smart *et.al.*(2001). The present study was initiated to standardize the technologies for producing true-to-type plants of guava in short period of time via soft wood cuttings with application of different concentrations of indole-3-butyric acid.

II. MATERIALS AND METHODS

The research studies were carried out under Net house condition, during the year 2013-14 and 2014-15.

Filling of poly bags: Before the cutting planting the 6 X 10 cm poly bags will be filled with different growing media. Six hundred (600) poly bags of each growing media should be filled i.e. 600 bags of soil, 600 bags of silt, 600 bags of soil + silt (1:1 ratio), 600 bags of Soil + Silt + FYM (1:1:1 ratio) and 600 bags of Soil + Silt + Vermicompost (1:1:1 ratio).

Preparation of cutting: After the filling of growing media in poly bags, the hard wood and semi hardwood cuttings of uniform size having 4-5 functional bud will be taken from healthy plants of Guava variety Allhabad safeda from one year matured shoots planted at the fruit research station Imaliya and gest house no- 2 guava orchard.

Preparation of Growth regulator solution: The weighing of IBA will be done with the help of electronic balance. The requisite quantity 1g, 2g and 3g (1000 ppm, 2000 ppm and 3000 mg) will be weighed separately and transferred in to flask and then initially the IBA sample will dissolve in 10 ml ethyl alcohol (90%) and make the volume 1000 ml with mixing 990ml distil water. By this process we will find the 1000, 2000 and 3000 ppm IBA solution.

Application of growth regulators: The fresh basal end cut of the cutting about 2.5 cm will be dipped in proposed hormonal solution for about 5 second and then after we will let in shade, so

that the cutting could absorb the hormone and best result should be obtained. After this process the cutting will be planted in poly bags.

Planting: The cutting about 0.75 to 1 cm thick diameter will be taken and planted in poly bags with 2-3 functional buds below the ground. Before the planting the hole should be done from planting place with the help of stick for preventing the buds to injuries.


Layout system of experiment was Factorial RBD replicated three times making 40 number of experimental units. Data on success percentage, length of sprouting (cm), total No. of leaves/cutting and number of roots taken after 120 days were collected during both years and were averaged and analyzed.

III. RESULTS AND DISCUSSION

Success percentage

The data (Table1) revealed that, the success percentage of cuttings in semi hard wood cuttings were recorded significantly highest 33.09% success with stem cuttings dipped in IBA @ 3000 ppm (G3), while, lowest 21.68% success of cuttings was recorded in G0 (control) in 1st year, 2nd year and pooled, respectively. In hard wood cuttings, stem cuttings dipped in IBA @ 3000 ppm (G3) were recorded significantly maximum 36.63% success of cuttings and it was lowest 23.90% in treatment G0 (control).

Usually higher dose of root promoting hormone inhibits the sprouting of initials (Urooba, *et.al.*, 2008), which happened truly in this experiment. Minimum success percentage (13%) was observed in cuttings kept as control. These results are in line with those of Manan *et al.* (2002) who reported that cuttings treated with IBA at 1000 ppm gave 37 percent success against control (17.5%). Abdullah *et al.* (2006) also observed that stem cuttings collected from mature stock plants gave the highest rooting percentage (60%) when treated with 0.4 percent IBA solution followed by two percent IBA.



Success percentage of cuttings

Length of Shoot (cm)

Stem cuttings dipped in IBA @ 3000 ppm (G_3) were recorded significantly maximum length of shoots 14.37 cm over the control (G_0) i.e. 7.76 cm length of shoots in semi hard wood cuttings. In hard wood cuttings, stem cuttings dipped in IBA @ 3000 ppm (G_3) were recorded significantly maximum 10.91 cm length of shoots and it was minimum 6.07 cm in treatment G_0 (control). Abdullah et al. (2006) also reported maximum shoot length (8.24 cm) soft wood cuttings treated with 1000 ppm IBA and minimum (3.83cm) in control treatment.

Total number of leaves/cutting

Application of 3000 ppm IBA (G_3) in dipping of stem cuttings were recorded significantly maximum 20.19 and minimum 11.62 total number of leaves cuttings⁻¹ were recorded under G_0 (control) in semi hard wood cuttings. In hard wood cuttings, stem cuttings dipping in the IBA 3000 ppm (G_3) were recorded significantly maximum 19.26 and it was minimum 10.75 leaves in treatment G_0 (control). Wahab *et al.* (2001) have also reported significantly higher number of leaves per cutting in 3000 ppm IAA at These results were also supported Rahman *et al.* (2004) who observed maximum number of leaves (10.20) in 1000 ppm IBA treatment.

Number of primary and secondary roots

Semi hard wood cuttings, (Table 1) results revealed that application of 3000 ppm IBA (G_3) in dipping of stem cuttings were recorded significantly maximum 26.58 and minimum 12.88 number of primary and secondary roots/cutting were observed in control (G_0). In hard wood cuttings, stem cuttings dipping in the IBA 3000 ppm (G_3) were recorded significantly maximum 24.10 and it was minimum 11.50 number of primary and secondary roots/cutting in treatment G_0 (control). Wahab et al. (2001) also reported that number of roots per cutting increased with higher IBA (4000ppm) against the lowest number of roots per cutting in 2000 ppm concentration. The present results also support the findings of Khatak et al. (1983). It was further observed that cuttings with more number of leaves produced more number of roots, due to fact that photosynthesis and other activities were carried out in leaves that caused more number of roots. Adventitious root formation is a key step for vegetative propagation comprising root induction, in which molecular and biochemical changes occur before any cytological event; root initiation when first anatomical modifications take place; and protrusion, corresponding to the emergence of root primordial (Berthon *et.al.*,1990 and Heloir *et.al.*,1996).). Lateral roots development in Arabidopsis provided a model for study of hormonal signals that regulated post embryonic organogenesis in higher plants (Zhang, H. and B.G. Forde.(2000) , Reddy, L.M. and R. N. Singh. (1988)). Lateral roots originated from pairs of pericycle cells, in several cell files positioned opposite the xylem pole, that initiated a series of asymmetric, transverse divisions to create 3 to 10 "short" daughter cells (Casimiro, I. *et. al.*,2001). These short daughter cells have undergone radial enlargement and subsequently divided periclinally to give rise to inner and outer cell layers. Further periclinal divisions resulted in formation of lateral root primordial (Bhalerao *et .al.*,2002).

Table 1 Effect of IBA on semi hard wood and hard wood cutting of guava

Semi hard wood cutting				
IBA (g)	Success percentage	Length of Shoot (cm)	Total number of leaves/cutting	Number of primary and secondary roots
G0	21.68	7.763	4.505	12.88
G1	25.38	9.716	4.831	16.68
G2	29.59	12.222	5.208	20.41
G3	33.09	14.369	5.441	26.58
Hard wood cutting				
G0	23.90	6.07	4.494	11.50
G1	27.01	7.15	4.790	15.00
G2	30.15	8.26	5.123	18.36
G3	36.63	10.91	5.333	24.10
Semi hard wood cutting		Hardwood cutting		
	SEm±	CD5%	SEm±	CD5%
1	0.51	1.45	0.84	2.40
2	0.23	0.68	0.34	0.98
3	0.05	0.14	0.045	0.13
4	0.258	0.739	0.235	0.673

IV. CONCLUSION

The study concludes that production of guava plants in net house condition proved to be the best for rapid and cheapest method multiplication of guava true-to-type plants. Significant results were obtained when guava nursery produced by semi hard wood and hard wood cuttings after application of 3000 ppm IBA. The plants produced by this technique will be true-to-type and can be planted in high density plane. These plants will bear earlier than the seedlings. The unique characters of a variety can be preserved through this technique. The technique was developed in this study is simpler, rapid, less labour intensive and economical, as root promoting hormones are required for root initiation. It is useful as compared to conventional method of propagation (grafting/budding) of guava because of higher success rate, independence of season and climate, small size of cuttings, use of juvenile shoot cuttings, disease free nature and production of large number of uniform true to mother type plants in a short period of time.

BIBLIOGRAPHY

- [1] Abdullah, T. M., M. A. Hussain and M. K. Bhuiyan. 2006. Clonal propagation of guava (*Psidium guajava* Linn.) by stem cutting from mature stock plants. J. Forestry Res. 17(4):301-304.
- [2] Bhalerao, R.P., J. Eklo, K. Ljung, A. Marchant, M. Bennett and G. Sandberg. 2002. Shoot derived auxin is essential for early lateral root emergence in Arabidopsis seedlings. Plant J. 29(3):325-332
- [3] Berthon, J.Y., B. Tahar and G.T. Boyer. 1990. Rooting phases of shoots of *Sequoiadendron giganteum* *in vitro* and their requirements. Plant Physiol. Biol. 28: 631-638.
- [4] Casimiro, I., A. Marchant, R.P. Bhalerao, T. Beeckman, S. Dhooge, R. Swarup, N. Graham, D. Inze, G. Sandberg, P.J. Casero and M. J. Bennett. 2001. Auxin transport promotes Arabidopsis lateral root initiation. Plant Cell. 13:843-852.
- [5] Heloir, M.C., C. Kevers, J. F. Hausman and T. Gaspar. 1996. Changes in the concentrations of auxins and polyamines during rooting of *in vitro* propagated walnut shoots. Tree Physiol. 16: 515-519.

- [6] Jaiswal, V.S. and M.N. Amin. 1987. In vitro propagation of guava from shoot cultures of mature tree. J. Pl. Physiol. 130:7-12.
- [7] Khattak, M. S., H. Inayatullah and S. Khan, 1983. Propagation of guava (*Psidium guajava* L.) from semi-hardwood cuttings. Frontier J. Agri. Res.,8:87-90.
- [8] Manan, A., M.A. Khan, W. Ahmad and A.Sattar. 2002. Clonal propagation of guava (*Psidium guajava* L.). Int. J. Agri. Biol. 1560- 8530/2002/04-1-143-144
- [9] Maqbool, M. and D.A. Khan. 1973. Propagation of fruit plants, studies on the propagation of guava by stem cuttings. Pak. J. Agri., 4: 177-193.
- [10] Martin, F.W., C.W. Campbell and R.M. Roberte. 1987. Perennial edible fruits of the tropics: An inventory. U.S. Dept. Agr., Agr. Res. Ser., Agr. Handb. 642.
- [11] Morton, J. 1987. Guava. In: Fruits of Warm Climates. Julia, F.(ed.). p. 356-363.
- [12] Pathak, R.K. and R.Dwivedi. 1996. Nursery practices of subtropical fruits suited for wasteland. Presented in Nursery Technology Workshop "Nursery'96" NBRI, Lucknow, India
- [13] Rahman, N., Tehsinullah, G. Nabi and T. Jan. 2004. Effect of different growth-regulators and types of cuttings on rooting of guava (*Psidium guajava* L.). Quarterly Sci. Vision. 9(1-2):1-5.
- [14] Reddy, L.M. and R. N. Singh. 1988. Efficacy of plastic house in propagation of guava (*Psidium guajava* L.)through hard wood cuttings. Indian J. Agri. Sci. 58:81-82
- [15] Samson, J.A. 1986. Tropical Fruits. 2nd Ed., Tropical Agriculture Series, Longman Scientific & Technical, Longman Inc., New York.
- [16] Singh, G., A.K. Singh and A. Verma. 2000. Economic evaluation of crop regulation treatments in guava (*Psidium guajava* L.), Indian J. Agric. Sci. 70:226-230
- [17] Singh, G., S. Gupta, R. Mishra and G.P. Singh. 2005. Wedge Grafting in Guava- A Novel Vegetative Propagation Technique. CISH, Lucknow. p.12.
- [18] Smart, D. R., L. Kocsis, M. A. Walker and C. Stockert. 2001. Dormant buds and adventitious root formation by Vitis and other woody plants, J. Pl. Growth Regulation. 21:296-314.
- [19] Steel, R.G.D., J.H. Torrie and D.A. Dickey. 2008. Principles and Procedures of Statistics. A biometrical Approach 3rd Ed. McGraw Hill Book Co., New York. U.S.A
- [20] Urooba, P., A. Khan, R. Javed and J. Zeb. 2008. Production of constraints of guava in district Kohat. Sarhad J. Agric. 24(3):449-554.
- [21] Wahab, F., G. Nabi, N. Ali and M. Shah. 2001. Rooting response of semi hardwood cuttings of guava (*Psidium guajava* L.) to various concentrations of different Auxins. J. Biol. Sci. 1(4):184-187.
- [22] Wilson, C.W. 1980. Guava. In: Tropical and Sub-tropical Fruits: Composition, Properties and Uses. Nagy, 5 and P.E. Shaw (eds.), AVI, Westport, CT.
- [23] Zhang, H. and B.G. Forde. 2000. Regulation of Arabidopsis root development by nitrate availability. J. Exp. Bot. 51: 51-59.

