



**A CASE OF**

**TRYPANOSOMOSIS IN A CAPTIVE ASIATIC WILD DOG (*Cuon alpinus*).**

**Boon Allwin<sup>1</sup>, Thirumurugan R<sup>2</sup>, Palanivelrajan, M<sup>3</sup>, Senthil N.R<sup>4</sup>, Vairamuthu. S<sup>5</sup>,  
Jayathangaraj .M.G <sup>6</sup>**

<sup>1</sup>Ph.D Scholar, Dept. of Wildlife Sciences, Madras Veterinary College, Chennai-07

<sup>2</sup>Arignar Anna Zoological park, Vandalur, Chennai-51

<sup>3</sup>Assistant professor, Dept. of Wildlife Sciences, Madras Veterinary College, Chennai-07

<sup>4</sup>Assistant professor, Centralized Clinical Laboratory, Madras Veterinary College, Chennai-07

<sup>5</sup>Professor and Head, Centralized Clinical Laboratory, Madras Veterinary College, Chennai-07

<sup>6</sup>Professor and Head, Dept. of Wildlife Sciences, Madras Veterinary College, Chennai-07

---

**ABSTRACT**

*Trypanosoma is an extra erythrocytic hemoprotozoan parasite, transmitted by biting flies and infected meat, causing fever, corneal opacity, anaemia. A male captive wild dog of Arignar Anna Zoological Park, Vandalur aged 3 years had a complaint of anorexia, lethargy, edema of forehead, staggering gait and reduced activity levels in its enclosure. Wet film revealed the presence of numerous motile Trypanosoma sp. Haematological parameters showed anaemia and mild neutrophilia. Serum biochemistry was found to be altered. The treatment done was administration of Triquin to the ailing animal. The wild dog was housed in the in-patient ward for continuous monitoring. No Trypanosomes could be detected in the wet film and stained smears taken subsequently. After a week, the wild dog was shifted to its enclosure and is doing well.*

**Key words: Wild dog, Trypanosomiasis, Haematology, Biochemistry, treatment**

---

**I. INTRODUCTION**

Trypanosomiasis is a hemoprotozoan disease of domestic and wild animals. Trypanosoma is an extra erythrocytic hemoprotozoan parasite, transmitted by biting flies and infected meat, causing fever, corneal opacity, anaemia, and myocarditis. Wild dog or Dhole is an endangered canid protected under Wildlife Protection Act, 1972 and a difficult species to maintain in captivity. All species of *Trypanosoma*, with the exception of some strains of *T. vivax* which produce a hyper acute and acute infection, characterized by high parasitaemia, fever, severe anemia and hemorrhages on the mucosal and serosal surfaces (Urquhart *et al.* 2002). Trypomastigote form of trypanosome enters host cells soon after infection, multiplies sub clinically and escapes the immune system and spread throughout the body primarily within macrophages. Parasitaemia develops within a few day and peaks 2 to 3 weeks post infection, coinciding with clinical disease (Barr *et al.* 1991). Anemia is a cardinal feature of the disease

in which red blood cells are removed from the circulation by the expelled mononuclear phagocytic system. Later, in infection of several months duration, when the parasitaemia become low and intermittent, anemia may resolve to a variable degree (Urquhart *et al.* 2002). There are a number of effective trypanosomacidal agents for dogs including suramin, quinapyramine and diminazene but single dose of diminazene aceturate is effective in eliminating the natural trypanosomiasis infection in canine (Rani and Suresh 2007).

## II. MATERIALS AND METHODS

A male captive wild dog aged 3 years weighing 15 kg, of Arignar Anna Zoological Park, Vandalur was presented with a complaint of anorexia, lethargy, edema of forehead, staggering gait and reduced activity in enclosure. The wild dog was chemically immobilized with xylazine @ 1mg/kg and ketamine @ 10 mg/kg and taken to the zoo veterinary hospital for detailed examination and treatment. Physical examination revealed high rise of temperature (40.8°C), pale mucous membrane, bilateral lacrimation, and generalized debility. On thoracic auscultation, exaggerated breath sounds on both sides were observed. The wet film, peripheral blood smear, whole blood and serum were collected. Hematology examination was done in Auto-haemoanalyser (BC-Vet 2800) and serum biochemistry in A15 auto analyser (Biosystems).

## III. RESULTS

. Wet film revealed numerous motile organisms that were suggestive of *Tryps* sp., the blood smear stained with Leishman-Giemsa stain revealed as many as 4-7 trypanosomes per field which is indicative of severe infection (Figure 1). Haematological parameters and Serum biochemistry was found to be altered (Table 1, 2&3). No abnormality was detected on thoracic radiography.

The ailing animal was treated with Triquin<sup>R</sup> (@ 0.025mg /kg BW) S/C single dose, plasma volume expanders, vitamins, I/V fluids and amino acids. The wild dog was housed in the in-patient ward for continuous monitoring. No Trypanosome could be detected in the wet film and smears taken subsequently. After a week, the wild dog was shifted to its enclosure and is doing well.

## IV. DISSCUSSION

There is no published literature available regarding the prevalence of trypanosomiasis in wild dog in India. Clinical signs observed in these wild dogs were in agreement with the findings reported by Rani and Suresh (2007), who reported *T. evansi* organism in peripheral blood with a history of inappetance, dullness and persistent fever since five days but the same authors also observed bilateral corneal opacity which is a characteristic finding in chronic trypanosomiasis also reported by Thirunavukkarasu *et al.* (2004). The fever characterized by high temperature might be due to the effects of toxic metabolites produced by dying trypanosomes (Tizard *et al.*, 1978).

The anaemic changes encountered are attributable to extravascular destruction of RBC's which may be through the process of erthro phagocytosis or metabolic product or from liberation of toxins by the parasites, hemodilution and depression of erythropoiesis. There was a marked decrease in RBC, Hb, and PCV and total WBC count remained unaltered as similar to findings observed by Hellebrekers & Slappendel, (1982) and Sandoval *et al*, (1994).

There were some alterations in the serum biochemistry that was in agreement with Lushaikyaa *et al.*, (2011). Amongst this creatinine, were found to be elevated and blood glucose was found to be decreased. The metabolic toxins liberated by the organisms may be the reason for low blood glucose level and glycogen reserve due to hepatic changes. Additionally, trypanosomes utilize a large amount of glucose to sustain their viability. An increase in ALT indicating hepatic degeneration, coinciding with the findings in T.evansi infection in dogs. There was a decrease in the albumin: globulin ratio. The fall in albumin levels was secondary to hyperglobulinemia as a compensatory mechanism for maintenance of normal blood viscosity increased by globulin levels.

In conclusion, clinical signs and symptoms along with the microscopic examination of *Trypanosoma* organism through peripheral blood smear suggested the case of trypanosomiasis. Moreover, it was also concluded that single dose of Triquin @ 0.025 mg/kg body weight successfully treated the wild dog with trypanosomiasis.

## V. ACKNOWLEDGEMENTS


The authors sincerely thank the Dean, Faculty of basic sciences, Madras Veterinary College, Dr. S. A. Asokan for the facilities provided and the Director, Arignar Anna Zoological park, Vandalur, Chennai-51

## BIBLIOGRAPHY

- Barr SC, Gossett KA, Klei TR. 1991. Clinical, clinicopathologic and parasitological observations of trypanosomiasis in dogs infected with North American *Trypanosoma cruzi* isolates. *Am J Vet Res.* 52 (1991) pp. 954-960.
- Hellebrekers LJ, Slappendel RJ. 1982. Trypanosomiasis in a dog imported in the Netherlands. *Vet. Q.* (1982) pp.182-186.
- Lushaikyaa A, David O, Rowland ISA, Sackey A KB. 2011. Hematological and serum biochemical changes in gilts experimentally infected with *Trypanosoma brucei*. *Veterinarski Arhiv* 81 (2011) pp.597-609.
- Rani NL, Suresh K. Canine trypanosomiasis. *Ind Vet J.* 84 (2007) pp.186-187.
- Sandoval GL, Coppo NB, Negrette MS. (1994). Alterações bioquímicas e histopatológicas de um cão e ratos infectados com *Trypanosoma evansi*. *Hora Vet.*, 81 (1994) pp. 53-55.

- Satheesh Kumar S. Health assessment of captive wild dogs. Master of Veterinary Science thesis submitted to the Madras Veterinary College, Tamil Nadu Veterinary and Animal Sciences University. Tamil Nadu. India. 2009.
- Thirunavukkarasu PS, Rao VV, Srinivasan SR, Nambi AP, Dhanapalan P.. *Ind J Vet Med.* 24 (2004) pp. 117.
- Tizard IR, Nielsen KH, Seed JR, and Hall JE. Biologically active products from African trypanosomes. *Microbiol. Rev.* 42 (1978):pp. 661-681.
- Urquhart G.M., Armour J, Duncan J.L, Dunn A.M, Jennings F.W . *Veterinary Parasitology.* 2nd Ed. Blackwell Science Co. UK. (2002). pp.217.

**FIGURE 1**



**TABLES**

**Table 1. Haematological examination**

<b>Component</b>	<b>Recorded value</b>	<b>Normal range*</b>
<b>Hb ( g/dl)</b>	9.2	(16.69 ± 0.26)
<b>RBC*10<sup>6</sup></b>	4.92	(7.49 ± 0.05)
<b>PCV (%)</b>	26.7%	(46.88 ±1.31)
<b>WBC*10<sup>3</sup></b>	7.2	(7.91 ± 4.79)
<b>MCV</b>	54.43	62.50
<b>MCH</b>	29.0	28.0
<b>Anaemia-microcytic normochromic</b>		

**Table.2 Differential count**

<b>Cell</b>	<b>Recorded value</b>	<b>Normal range*</b>
<b>Neutrophils</b>	74	(72.63 ± 2.81)
<b>Lymphocytes</b>	20	(25.25 ± 2.76)
<b>Monocytes</b>	02	(00.75 ± 0.31)
<b>Eosinophils</b>	04	(01.38 ± 0.41)
<b>Basophils</b>		NIL

**Table 3. Serum Biochemistry**

<b>Parameters</b>	<b>Recorded value</b>	<b>Normal range*</b>
<b>BUN</b>	20.40	(20.56 ± 0.74)
<b>Creatinine</b>	2.87	(1.44 ± 0.04)
<b>Total protein (g/dl)</b>	7.03	(6.09 ± 0.30)
<b>Albumin (g/dl)</b>	3.01	(2.38 ± 0.22)
<b>Globulin (g/dl)</b>	4.02	(3.71 ± 0.33)
<b>Calcium (mg/dl)</b>	2.70	(9.10 ± 0.64)
<b>Phosphorus(mg/dl)</b>	6.07	(5.86 ± 0.48)
<b>Glucose</b>	33.99	(53.63 ± 1.28)
<b>ALT</b>	102	(19.76 ± 0.79)

Normal range \* (Satheesh Kumar 2009)